REPORT DOCUMENTATION PAGE OMB No. 0704-0188 Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS. 3. DATES COVERED (From - To) 1. REPORT DATE (DD-MM-YYYY) 2. REPORT TYPE View Graphs 6 May 2003 4. TITLE AND SUBTITLE 5a. CONTRACT NUMBER **5b. GRANT NUMBER** Structural Effects on the Physical Properties of Ionic Liquids 5c. PROGRAM ELEMENT NUMBER 6. AUTHOR(S) 5d. PROJECT NUMBER 2303 Greg Drake, et al. 5e. TASK NUMBER M2C8 5f. WORK UNIT NUMBER 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZATION REPORT NUMBER Air Force Research Laboratory (AFMC) AFRL-PR-ED-VG-2003-122 AFRL/PRSP 10 E. Saturn Blvd. Edwards AFB, CA 93524 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSOR/MONITOR'S ACRONYM(S) Air Force Research Laboratory (AFMC) 11. SPONSOR/MONITOR'S AFRL/PRS NUMBER(S) 5 Pollux Drive AFRL-PR-ED-VG-2003-122 Edwards AFB CA 93524-7048 12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited. 13. SUPPLEMENTARY NOTES 14. ABSTRACT 20030616 039 15. SUBJECT TERMS 16. SECURITY CLASSIFICATION OF: 17. LIMITATION 18. NUMBER 19a, NAME OF RESPONSIBLE OF ABSTRACT OF PAGES **PERSON** Sheila Benner

Α

c. THIS PAGE

Unclassified

b. ABSTRACT

Unclassified

a. REPORT

Unclassified

code)

(661) 275-5693

19b. TELEPHONE NUMBER (include area

Form Approved



MEMORANDUM FOR PRS (In-House Publication)

FROM: PROI (STINFO)

06 May 2003

SUBJECT: Authorization for Release of Technical Information, Control Number: AFRL-PR-ED-VG-2003-122

Greg Drake (AFRL/PRSP) et al., "Structural Effects on the Physical Properties of Ionic Liquids"

2003 AFOSR Molec Dynamics & Theo Chem Contr Mtg (San Diego, CA, no date provided) (Deadline: 19 May 2003) (Statement A)

Structural Effects on the Physical Properties of Ionic Liquids

Greg Drake and Tom Hawkins AFRL/PRSP Air Force Research Laboratory Edwards AFB, CA 93524

and

John Wilkes
Department of Chemistry
United States Air Force Academy
USAF Academy, CO 80840



Distribution Statement A: Public Release unrestricted








NOT

versus



Extended lattice

Cryolite Na₃AlF₆ m.p. nearly 1000 °C (Hall Process for Al production) Table salt Na⁺Cl⁻ m.p. = 804 °C Very high Eutectic of Li+Cl- and K+Cl- m.p. 355 °C

Molten salts are very hot!

Not commercially viable

Corrosion and energy issues

Giant lattice of miniature magnets stuck together







What are Ionic Liquids?

A class of salts consisting of cation/anion pair that has a very low melting point.

Definition of an ionic liquid is open to some debate amongst researchers in the area, but most in the area use one of two. (1) An ionic compound that melts below 100 °C (b.p. of H₂O). J. Wilkes, P. Wasserscheid, K. Seddon. (2) An ionic compound that has a melting point at or below ambient temperatures. These are often called RTILs (Room Temperature Ionic Liquids) T. Welton, R. Rogers.

But many of the salts fit both definitions and 2 is really a more specific class of (1).





Important factors affecting the physical properties of ionic liquids

1. Asymmetry of cation as well as anion


2. Packing efficiency


3. Charge delocalization in cationic/anionic species


4. "Sheer size" differentials














[CH₃CH₂NH₃⁺][NO₃-] m.p. 12 °C Ethylammonium nitrate


Serious issues...

- can be treacherous
- acidic
- -very hygroscropic







Some major shapes for organic based cations



1-methyl-3-alkyl-imidazolium



1-alkylpyridinium



Tetralkylphosphonium









The group of anions for ionic liquids is much larger and growing....


water soluble	water insoluble
CH,CO,-	PF_6
CF,CO,-	$[\mathbf{BR}_1\mathbf{R}_2\mathbf{R}_3\mathbf{R}_4]$
Cl-, Br-, I-	$[(CF_3SO_2)_2N]$
NO 3-	BF_4
BF4	$R-SO_3$ -
NO_2 -	
$[AICI_4, AI_2CI_7]$	

This list is not comprehensive but it covers the majority of what is out there. Typically R groups are n-alkyl groups





aluminum trichloride systems. More complex than originally thought as AICl₃ Significant efforts spent on 1-ethyl-3-methyl-imidazolium based systems and and CI have an equilibrium based on their respective concentrations.





"Strongly acidic"

"Acidic"

Wilkes, J. S. Green Chemistry 2002, 4, 3.








Fannin, A.; Floreani, D.; King, L.; Landers, J.; Piersma, B.; Stetch, D.; Vaughn, R.; Wilkes, J.; Williams, J. J. Phys. Chem. 1984, 88, 2614.









.Seddon, K.R.; Holbrey, J. D. Clean Products and Processes 1999, 1, 223. Rogers, R.; Seddon, K. (eds.) Ionic Liquids A.C.S. Symp. Ser. 818 2002 A.C.S Publ. Co. Wasserscheid, P.; Keim, W. Angew. Chem. Int. Ed. Engl. 2000, 39, 3772. Wasserscheid, P, Welton, T. (eds.) Ionic Liquids in Synthesis Wiley-VCH, FRG, 2003



Ionic Liquids melting points



\$	× \
(Z
	Z
	Н3С—
	


3-methyl-1-R-imidazolium

Substituent	Triflate (m.p.)	Bis(trifluorosulfonamide) m.p.
1-methyl	39	22
1-ethyl	6-	ę
1-butyl	16	4-
1-СН, ОСН, СН,	3 27	<-30(Tg)
$1-\mathrm{CH}_2^-\mathrm{CF}_3^-$	45	<-30(Tg)


R	•	
	†\ +\ -\	
2		$\widetilde{\mathbf{H}}_2$
J II	113~	

Bis(trifluorosulfonamide) m.p.	14	<-30(Tg)
Triflate (m.p.)	23	2
Substituent	1-ethyl	1-butyl

3-ethyl-1-R-imidazolium



1-ethyl-2-R-3-methyl-imidazolium



1-ethyl-3-R-5-methyl-imidazolium





8 00 8 5 6 8 5 6

8.4 8.8 3.9 4.2 0.98

34 (45) 52 (90) 54 (74)

\(mS/cm)

1(cP)

Bis(trifluoromethylsulfonamide)

-methyl

1-ethyl 1-butyl

Z X
H_3

3-methyl-1-R-imidazolium


1-CH₂OCH₂CH₃

-CH₂-CF₃

N/N	
$\langle z \rangle$	\exists
r () H
H,C	9

3-ethyl-1-R-imidazolium

Bis(trifluoromethylsulfonamide)	$\eta(cP)$	A(mS/cm)
1-ethvl	35 (43)	2 C) 2 X
I-cury.	33 (23)	200
1-butyl	48	4.1



Bis(trifluoromethylsulfonamide)	η(cP)	Λ(mS/cm)
3-methyl	37 (51)	(6.4)
3-ethyl	36	6.2

Red values in () are for corresponding CF3SO3- salt

1-ethyl-3-R-5-methyl-imidazolium


Viscosity and conductivity increase with increasing chain length but intramolecular hydrogen bonding can be important. Size and "charge" of anion also significant.

Bonhote, P.; Diaz, A.; Papageorgiou, N.; Kalyanasundaram, K.; Gratzel, M. Inorg. Chem. 1996, 35, 1168.





carborane anion in the formation of ionic liquids. Essentially no hydrogen bonding. The end extreme of non-coordinating anions has been achieved through the use of



Ionic liquid	m.p. °C
$[EMIM][HCB_{11}H_{11}]$	122
$[EMIM][1-CH_3-CB_{11}H_{11}]$	20
$[EMIM][1-CH_2CH_3-CB_{11}H_{11}]$	64
$[EMIM][1-CH_2CH_2CH_3-CB_{11}H_{11}]$	45
[EMIM][1-CH, CH, CH, CH, -CB, H, .]	49


-imidazolium icosahedral	rane [EMIM][1-prop-CB ₁₁ H ₁₁]
1-ethyl-3-methyl-imid	1-propyl-1-carborane





$-CF_3$	Salt	m.p.(°C)	m.p.(°C) DSC onset (°C)
	NTf ₂ -	L9-	395
N.	Tf0-	33	379
1-(3',3',3'-trifluoro-n-propyl)-3-n-butyl-1,2,4-triazolium			
CH_2F	Salt	m.p.(°C)	m.p.(°C) DSC onset (°C)
	NTf.	-70	359
	$\mathbf{BF_4}^-$	52	336

1-(2'-fluoroethyl)-3-n-heptyl-1,2,4-triazolium




1-(1H,1H,2H,2H-perfluoro-n-hexyl)-3-n-butyl-1,2,4-triazolium






Substituted ammonium salts R, N+X- Variations in melting point based on cation structure.




Tris-(n-propyl)-undecylammonium cation ClO_4 m.p. = 65 ${}^{0}C$ Br m.p. = 67° C




Tetra-n-pentylammonium cation

 ClO_4 m.p. = 118 $^{\circ}C$ Br m.p. = $101 \, ^{\circ}$ C



N-decyl-n-octyl-dimethylammonium cation Br. m.p. = RTIL, CIO₄· m.p. = RTIL



Br m.p. = 170° C, ClO₄ m.p. = 152° C N-tetradecyl-triethylammonium cation



Gordon, J. E.; SubbaRao, G. N. J. Amer. Chem. Soc. 1978, 100, 7445.





Substituted ammonium salts [R₄N⁺][X-]Recently work has been done by using more desirable anions.

Substituted Ammonium Salt	M.P.	Density	Viscosity	V
	(° C)	(g/cm^3)	(cb)	(Ω-1 cm²/mole)
$[(n-C_cH_{1,2})(CH_1)_2N^+][N(SO_2CF_3)_2]$		1.33	153	1.4
$[(\mathbf{n} - \mathbf{C}_7 \mathbf{H}_{15})(\mathbf{C}\mathbf{H}_3)_3 \mathbf{N}^+][\mathbf{N}(\mathbf{SO}, \mathbf{CF}_3)_7]$		1.28	153	1.4
$[(\mathbf{n} - \mathbf{C}_{\mathbf{c}} \mathbf{H}_{17})(\mathbf{C} \mathbf{H}_{3})_{\mathbf{i}} \mathbf{N}^{+}][\mathbf{N}(\mathbf{SO}, \mathbf{CF}_{3})_{\mathbf{i}}]$		1.27	181	1.3
[(n-C,H,1)(CH,CH,),N+][N(SO,CF1),-]		1.27	167	2.5
[(n-C,H,s)(CH,CH,),N+][N(SO,CF,),-]		1.26	75	1.9
$[(n-C_0H_{1,7})(CH_1CH_2),N^+][N(SO,CF_3),-]$		1.25	202	1.3
$[(n-C_cH_{1,2})(n-C_dH_0)_3N^+][N(SO,CF_3)_7]$		1.15	595	8.0
$[(n-C_7H_{1z})(n-C_4H_0)_3N^+][N(SO,CF_3)_7]$		1.17	909	8.0
$[(n-C_0H_{1,7})(n-C_dH_0)_3N^+][N(SO_2CF_3)_7]$	-63	1.12	574	0.7
[(n-C,H,z)(Et),(ipr),N+[[N(SO,CF3),-]		1.27	362	1.2
$[(n-C_8H_{17})(n-C_4H_9)_3N^+][OSO_2CF_3]$		1.02	2030	0.07

-most have very low glass points

-densities decrease as expected


-viscosity increases dramatically with increasing alkyl length

-conductivity decreases with cation size (mobility issue)

Sun, J.; Forsyth, M.; MacFarlane, D. R. J. Phys. Chem. B 1998, 102, 8858.








1-n-butyl-4-amino-1,2,4-triazolium bromide

1-substituted 4AT salts	m.p. (°C)	m.p. (°C) dec. onset (°C)	density (g/cm ³)
1-ethyl	63°	110	1.69
1-n-propyl	₀ 09	120	1.56
1-isopropyl	°06	110	1.60
1-butyl	48°	130	1.46
1-n-pentyl	54°	130	1.37
1-n-hexyl	°97	120	1.34
1-n-heptyl	940	120	1.30
1-n-octyl	80°	135	1.27
1-n-nonyl	81°	140	1.26
1-n-decyl	°06	135	1.23

*Drake, G. W.; Hawkins, T. W.; Tollison, K.; Hall, L.; Vij, A. 2003 manuscript in progress.







Extensive hydrogen bonding in 1-n-propyl-4-amino-1,2,4-triazolium bromide

*Drake, G. W.; Hawkins, T. W.; Tollison, K.; Hall, L.; Vij, A. 2003 manuscript in progress.









1-dodecyl-3-methylimidazolium hexafluorophosphate*

1-hexyl-4-amino-1,2,4-triazolium bromide#

*Gordon, C. M.; Holbrey, J. D.; Kennedy, A. R.; Seddon, K. R. J. Mater. Chem. 1998, 8, 2627.

*Drake, G. W.; Hawkins, T. W.; Tollison, K.; Hall, L.; Vij, A. 2003 manuscript in progress.





Summary and Conclusions

- Overall cation symmetry or lack thereof dramatically affects the physical properties of ionic liquids.
- Inter- as well as intra- molecular interactions especially hydrogen bonding are very important.
- Conductivity and viscosity are indirectly related, and both are significantly affected by the size and charge distribution of the cation and/or anion.
- New classes of ionic liquids are appearing and the field has tremendous promise for new and exciting breakthroughs.





Acknowledgements:

- Mike Berman (AFOSR)

- Kerri Tollison, Leslie Hall, Greg Kaplan, Ashwani Vij (coworkers)

- Wayne Kalliomaa and Ronald Channell (AFRL/PRSP)

- Mike Huggins (AFRL/PRS)